arXiv:hep-ph/9804396v1 24 Apr 1998

THE WEAK ISOSPIN SPACE

Gunn A. Quznetsov

February 7, 2008

Abstract

The Clifford pentad of the 4X4 matrices defines the 5-dimensional space. Each weak isospin transformation divides an electron on two components, which scatter in the 2-dimensional subspace and which indiscernible in the orthogonal 3-dimensional subspace.

PACS 12.15.-y

In the weak isospin theory we have got the following entities:
the right electron state vector e_R,
the left electron state vector e_L,
the electron state vector e ($e = \begin{bmatrix} e_R \\ e_L \end{bmatrix}$),
the left neutrino state vector ν_L,
the zero vector right neutrino ν_R.
the unitary 2×2 matrix U of the isospin transformation.($\det(U) = 1$).
This matrix acts on the vectors of the kind:$\begin{bmatrix} \nu_L \\ e_L \end{bmatrix}$.
Therefore, in this theory: if

$$U = \begin{bmatrix} u_{1,1} & u_{1,2} \\ u_{2,1} & u_{2,2} \end{bmatrix}$$

then the matrix

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & u_{1,1} & 0 & u_{1,2} \\ 0 & 0 & 1 & 0 \\ 0 & u_{2,1} & 0 & u_{2,2} \end{bmatrix}$$

operates on the vector

$$\begin{bmatrix} e_R \\ e_L \\ \nu_R \\ \nu_L \end{bmatrix}.$$

Because e_R, e_L, ν_R, ν_L are the two-component vectors then

$$\begin{bmatrix} e_R \\ e_L \\ \nu_R \\ \nu_L \end{bmatrix} \text{ is } \begin{bmatrix} e_{R1} \\ e_{R2} \\ e_{L1} \\ e_{L2} \\ 0 \\ 0 \\ \nu_{L1} \\ \nu_{L2} \end{bmatrix}$$

and

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & u_{1,1} & 0 & u_{1,2} \\ 0 & 0 & 1 & 0 \\ 0 & u_{2,1} & 0 & u_{2,2} \end{bmatrix} \text{ is } \underline{U} = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & u_{1,1} & 0 & 0 & 0 & u_{1,2} & 0 \\ 0 & 0 & 0 & u_{1,1} & 0 & 0 & 0 & u_{1,2} \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & u_{2,1} & 0 & 0 & 0 & u_{2,2} & 0 \\ 0 & 0 & 0 & u_{2,1} & 0 & 0 & 0 & u_{2,2} \end{bmatrix}.$$

This matrix has got eight orthogonal normalized eigenvectors of kind:

$$\underline{s_1} = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \underline{s_2} = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \underline{s_3} = \begin{bmatrix} 0 \\ 0 \\ \varpi \\ 0 \\ 0 \\ 0 \\ \chi \\ 0 \end{bmatrix}, \underline{s_4} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ \chi^* \\ 0 \\ 0 \\ 0 \\ -\varpi^* \end{bmatrix},$$

$$\underline{s_5} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \underline{s_6} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \underline{s_7} = \begin{bmatrix} 0 \\ 0 \\ \chi^* \\ 0 \\ 0 \\ 0 \\ -\varpi^* \\ 0 \end{bmatrix}, \underline{s_8} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ \varpi \\ 0 \\ 0 \\ 0 \\ \chi \end{bmatrix}.$$

The corresponding eigenvalues are: 1, 1, $\exp(i \cdot \lambda)$, $\exp(i \cdot \lambda)$, 1,1, $\exp(-i \cdot \lambda)$, $\exp(-i \cdot \lambda)$.

These vectors constitute the orthogonal basis in this 8-dimensional space.

Let

$$\underline{\gamma^0} = \begin{bmatrix} \gamma^0 & O \\ O & \gamma^0 \end{bmatrix}$$

if O is zero 4×4 matrix and

$$\gamma^0 = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix},$$

and $\underline{\beta^4} = \begin{bmatrix} \beta^4 & O \\ O & \beta^4 \end{bmatrix}$ if

$$\beta^4 = \begin{bmatrix} 0 & 0 & i & 0 \\ 0 & 0 & 0 & i \\ -i & 0 & 0 & 0 \\ 0 & -i & 0 & 0 \end{bmatrix}.$$

The vectors $\begin{bmatrix} e_{R1} \\ e_{R2} \\ e_{L1} \\ e_{L2} \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$, $\begin{bmatrix} e_{R1} \\ e_{R2} \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$, $\begin{bmatrix} 0 \\ 0 \\ e_{L1} \\ e_{L2} \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$ correspond to the state vectors e, e_R and e_L resp.

In this case [1] $\underline{e}^\dagger \cdot \underline{\gamma^0} \cdot \underline{e} = J_0$, $\underline{e}^\dagger \cdot \underline{\beta^4} \cdot \underline{e} = J_4$, $J_0 = \underline{e}^\dagger \cdot \underline{e} \cdot V_0$, $J_4 = \underline{e}^\dagger \cdot \underline{e} \cdot V_4$.

For the vector e the numbers k_3, k_4, k_7, k_8 exist, for which: $\underline{e} = (e_{R1} \cdot \underline{s_1} + e_{R2} \cdot \underline{s_2}) + (k_3 \cdot \underline{s_3} + k_4 \cdot \underline{s_4}) + (k_7 \cdot \underline{s_7} + k_8 \cdot \underline{s_8})$.

Here $\underline{e_R} = (e_{R1} \cdot \underline{s_1} + e_{R2} \cdot \underline{s_2})$. If $\underline{e_{La}} = (k_3 \cdot \underline{s_3} + k_4 \cdot \underline{s_4})$ and $\underline{e_{Lb}} = (k_7 \cdot \underline{s_7} + k_8 \cdot \underline{s_8})$ then $\underline{U} \cdot \underline{e_{La}} = \exp(i \cdot \lambda) \cdot \underline{e_{La}}$ and $\underline{U} \cdot \underline{e_{Lb}} = \exp(-i \cdot \lambda) \cdot \underline{e_{Lb}}$.

Let for all k $(1 \leq k \leq 8)$: $\underline{h_k} = \underline{\gamma^0} \cdot \underline{s_k}$. The vectors $\underline{h_k}$ constitute the orthogonal basis, too. And the numbers q_3, q_4, q_7, q_8 exist, for which: $\underline{e_R} = (q_3 \cdot \underline{h_3} + q_4 \cdot \underline{h_4}) + (q_7 \cdot \underline{h_7} + q_8 \cdot \underline{h_8})$.

Let $\underline{e_{Ra}} = (q_3 \cdot \underline{h_3} + q_4 \cdot \underline{h_4})$, $\underline{e_{Rb}} = (q_7 \cdot \underline{h_7} + q_8 \cdot \underline{h_8})$, $\underline{e_a} = \underline{e_{Ra}} + \underline{e_{La}}$ and $\underline{e_b} = \underline{e_{Rb}} + \underline{e_{Lb}}$.

Let $\underline{e_a}^\dagger \cdot \underline{\gamma^0} \cdot \underline{e_a} = J_{0a}$, $\underline{e_a}^\dagger \cdot \underline{\beta^4} \cdot \underline{e_a} = J_{4a}$, $J_{0a} = \underline{e_a}^\dagger \cdot \underline{e_a} \cdot V_{0a}$, $J_{4a} = \underline{e_a}^\dagger \cdot \underline{e_a} \cdot V_{4a}$, $\underline{e_b}^\dagger \cdot \underline{\gamma^0} \cdot \underline{e_b} = J_{0b}$, $\underline{e_b}^\dagger \cdot \underline{\beta^4} \cdot \underline{e_b} = J_{4b}$, $J_{0b} = \underline{e_b}^\dagger \cdot \underline{e_b} \cdot V_{0b}$, $J_{4b} = \underline{e_b}^\dagger \cdot \underline{e_b} \cdot V_{4b}$.

In this case: $J_0 = J_{0a} + J_{0b}$, $J_4 = J_{4a} + J_{4b}$.

Let $\left(\underline{U} \cdot \underline{e_a}\right)^\dagger \cdot \underline{\gamma^0} \cdot \left(\underline{U} \cdot \underline{e_a}\right) = J'_{0a}$, $\left(\underline{U} \cdot \underline{e_a}\right)^\dagger \cdot \underline{\beta^4} \cdot \underline{\left(\underline{U} \cdot \underline{e_a}\right)} = J'_{4a}$, $J'_{0a} = \left(\underline{U} \cdot \underline{e_a}\right)^\dagger \cdot \left(\underline{U} \cdot \underline{e_a}\right) \cdot V'_{0a}$, $J'_{4a} = \left(\underline{U} \cdot \underline{e_a}\right)^\dagger \cdot \left(\underline{U} \cdot \underline{e_a}\right) \cdot V'_{4a}$,

$\left(\underline{U}\cdot\underline{e_b}\right)^\dagger\cdot\underline{\gamma}^0\cdot\left(\underline{U}\cdot\underline{e_b}\right)=J'_{0b}$, $\left(\underline{U}\cdot\underline{e_b}\right)^\dagger\cdot\underline{\beta}^4\cdot\left(\underline{U}\cdot\underline{e_b}\right)=J'_{4b}$, $J'_{0b}=\left(\underline{U}\cdot\underline{e_b}\right)^\dagger\cdot\left(\underline{U}\cdot\underline{e_b}\right)\cdot V'_{0b}$, $J'_{4b}=\left(\underline{U}\cdot\underline{e_b}\right)^\dagger\cdot\left(\underline{U}\cdot\underline{e_b}\right)\cdot V'_{4b}$.

In this case:

$$V'_{0a}=V_{0a}\cdot\cos\left(\lambda\right)+V_{4a}\cdot\sin\left(\lambda\right),$$

$$V'_{4a}=V_{4a}\cdot\cos\left(\lambda\right)-V_{0a}\cdot\sin\left(\lambda\right);$$

$$V'_{0b}=V_{0b}\cdot\cos\left(\lambda\right)-V_{4b}\cdot\sin\left(\lambda\right),$$

$$V'_{4b}=V_{4b}\cdot\cos\left(\lambda\right)+V_{0b}\cdot\sin\left(\lambda\right).$$

Hence, every isospin transformation divides a electron on two components, which scatter on the angle $2\cdot\lambda$ in the space of (J_0, J_4). These components are indiscernible in the space of (j_1, j_2, j_3) [2].

My other addresses are:
gunn.q@usa.net
quznets@geocities.com

References

[1] xxx.lanl.gov/physics/9803047 p.3

[2] Item, p.2, p4